Inadequate distribution of dietary protein and muscle mass in older adults

Distribuição inadequada de proteínas nas refeições e massa muscular de idosos

Luciana Bronzi de Souza^a , Karine Anusca Martins^a , Rafael Aiello Bomfim^b

^aUniversidade Federal de Goiás – Goiânia (GO), Brazil.

^bUniversidade Federal de Mato Grosso do Sul – Campo Grande (MS), Brazil.

Address for correspondence

Luciana Bronzi de Souza – Universidade Federal de Goiás – Rua 227 Qd. 68 s/n – Setor Leste Universitário – CEP: 74605-080 – Goiânia (GO), Brasil. E-mail: lucianabronzi@ufg.br

Received on: 5/25/2021

Accepted on: 9/24/2021

How to cite this article: Souza LB, Martins KA, Bomfim RA. Inadequate distribution of dietary protein and muscle mass in older adults. *Geriatr Gerontol Aging*. 2022;16:e0220001. <https://doi.org/10.53886/gga.e0220001>

Abstract

Objective: To assess protein intake, protein distribution across meals, and a possible association between protein distribution and muscle mass in older adults.

Methods: This cross-sectional study included 47 older adults attending a community center in Campo Grande, MS, Brazil. A sociodemographic questionnaire was used to characterize the population. Dietary intake data were collected using two 24-hour dietary recalls on nonconsecutive days. Anthropometric measures, such as weight, height, and body circumferences, and measures of handgrip strength were collected. Skeletal muscle mass was calculated from the anthropometric measures. Descriptive statistics were used to derive means and standard deviations. Analysis of variance and Tukey tests were performed, with a significance level of 5% ($p < 0.05$).

Results: The older adults, mostly Caucasian, female, physically active, at risk of malnutrition, and with a low level of education, had an energy intake of 1,305.28 kcal/day. Protein intake was 65.19 g/day, with a mean of 0.93 g/kg/day. Lunch was the meal with the highest protein intake (33.85 ± 2.47 g). The lowest protein intake was observed at breakfast (8.56 ± 0.82 g). All meals were below the recommendation of 40 g of protein/meal. Protein intake at breakfast was associated with calf circumference ($p = 0.04$).

Conclusions: The older adults consumed a low amount of total and per-meal protein. Breakfast showed the lowest protein intake, and this was associated with calf circumference in the study sample.

Keywords: aged; aging; sarcopenia; food consumption; frailty.

Resumo

Objetivo: Avaliar o consumo proteico, sua distribuição nas principais refeições e sua possível associação com a massa muscular de idosos.

Metodologia: Estudo transversal realizado com 47 idosos frequentadores de um centro de convivência em Campo Grande (MS). Aplicou-se um questionário sociodemográfico e dois recordatórios de 24 horas, em dias alternados, para a avaliação do consumo alimentar, e realizou-se avaliação antropométrica por meio da aferição de medidas como peso, estatura, circunferências corporais e a força de pressão palmar. Com as medidas antropométricas, foi calculada a quantidade de massa muscular esquelética. Realizou-se estatística descritiva e, para as análises de associação, aplicaram-se os testes de variância (ANOVA) e de Tukey, com significância estatística quando $p < 0,05$.

Resultados: Os idosos, majoritariamente caucasianos, do sexo feminino, fisicamente ativos, em risco de desnutrição e com escolaridade reduzida apresentaram consumo energético de 1.305,28 kcal/dia e consumo proteico 65,19 g/dia, sendo 0,93 g de proteína/kg/dia. O almoço foi a refeição com maior aporte proteico ($33,85 \pm 2,47$ g). O menor consumo proteico foi observado no café da manhã ($8,56 \pm 0,82$ g). Todas as refeições ficaram abaixo da recomendação de 40 g de proteína/refeição. O consumo proteico no café da manhã associou-se à circunferência da panturrilha ($p = 0.04$).

Conclusões: Os idosos consumiram baixa quantidade de proteínas totais e nas principais refeições. O café da manhã foi a refeição com menor aporte proteico e este associou-se à circunferência da panturrilha em idosos da amostra.

Palavras-chave: idoso; envelhecimento; sarcopenia; consumo de alimentos; fragilidade.

This article is published in Open Access under the Creative Commons Attribution license, which allows use, distribution, and reproduction in any medium, without restrictions, as long as the original work is correctly cited.

INTRODUCTION

The older adult population has increased considerably in recent decades worldwide. In 2020, 9.8% of the Brazilian population consisted of individuals aged 65 years or over, and that number is expected to grow to more than 20.0% by 2046.¹ Not only is the number of older adults increasing but also their longevity. Life expectancy in Brazil was 76.6 years in 2019 and is estimated to reach 81.2 years by 2060.¹

Aging is characterized by changes in body composition, including increased adiposity and decreased lean mass, especially skeletal muscle mass (SMM).² The loss of muscle mass, muscle strength, and physical function that progresses with aging, known as sarcopenia, is a geriatric syndrome that causes impaired motor function, compromised autonomy, poorer quality of life, increased risk of falls and fractures, difficulty performing daily activities, increased risk and cost of hospitalization, and increased risk of morbidity and mortality.^{2,3} Recently, the revised consensus on definition and diagnosis of sarcopenia of the European Working Group on Sarcopenia in Older People defined low muscle strength as an independent indicator of probable sarcopenia.³

Medical literature data highlight the importance of protein intake in preventing and treating sarcopenia and in maintaining muscle health in old age.^{4,5} Protein intake stimulates SMM synthesis and inhibits its breakdown, resulting in a positive protein balance and muscle mass gain.² Also, not only is the total amount of protein consumed associated with muscle mass in older adults but also its distribution across meals, so that muscle protein synthesis (MPS) is stimulated several times per day.⁶ A protein intake of 40 g in each of the main meals is believed to be required to stimulate MPS in older adults.⁷ The current daily protein intake recommendation (0.8 g/kg/d)⁸ seems to be insufficient to achieve the ideal level of MPS stimulation, especially in older adults,⁴ and to guarantee an adequate protein distribution across meals.

Because remaining independent and functional is a key condition for the health of the aging individual, investigating effective factors to mitigate sarcopenia is crucial. However, little is known about this topic to date. Thus, given the lack of more detailed information, this study aimed to evaluate protein intake, protein distribution across the main meals, and a possible association of protein distribution with muscle mass in older adults.

METHODS

Study design, site, and sample

This population-based, descriptive, cross-sectional study was conducted in a community center for older adults in Campo

Grande, Mato Grosso do Sul, Brazil. Data were collected from June to October 2017.

The sample consisted of both men and women, aged 60 years or over, who regularly attended the community center. Individuals with disabling diseases (paralysis, stroke, difficulty walking), physical limitations (amputations), and those who did not fully comply with the research protocol were excluded.

Convenience sampling was used in this study. All 284 older adults registered in the community center were invited to participate; however, only 47 were included and completed the protocol.

All data were collected by a research team consisting of previously trained Nutrition students and nutritionists who were responsible for questionnaires, interviews, and nutritional assessments.

Anthropometric assessment

Anthropometric measurements were based on standardized procedures.⁹ All measurements were taken twice with the participant wearing minimal clothing. If the difference was greater than 0.1 kg or 0.1 m, a third measurement was performed, and the average of the two closest measurements was used in the analysis. Anthropometric assessment consisted of measuring weight (kg), height (m), waist circumference (WC), arm circumference, and calf circumference (CC) (cm). Circumferences were measured with an inelastic measuring tape, with a precision of 1 mm.

Body mass index (BMI) was obtained by the ratio between body weight and squared height ($BMI = \text{weight} / \text{height}^2$) and classified according to the levels proposed by Lipschitz.¹⁰ WC was assessed with the cutoff points suggested by the National Cholesterol Education Program – Adult Treatment Panel III,¹¹ whose defining levels for greatly increased risk for cardiovascular disease are > 102 cm for men and > 88 cm for women. CC was classified according to the levels used in the Mini-Nutritional Assessment (MNA®), which considers CC ≥ 31 cm to be adequate for both men and women. Lower levels are indicative of risk of malnutrition in older adults.

In addition to anthropometric measurements, the MNA® score was used to identify older patients at risk of malnutrition with the following cutoff points: adequate, ≥ 24; risk of malnutrition, 17 to 23.5; malnourished, < 17.

Muscle mass

Based on anthropometric data, the equation proposed by Lee et al.¹² was used to estimate appendicular skeletal muscle mass (ASM): $ASM = (0.244 \times \text{weight}) + (7.8 \times \text{height}) + (6.6 \times \text{gender}) - (0.098 \times \text{age}) + (\text{race} - 3.3)$. Zero (0) was

used for women and 1 for men; 0 for White or Hispanic; 1.4 for Afro-descendant; and -1.2 for Asian. This equation showed a strong agreement with dual-energy X-ray absorptiometry (DXA) and with the prevalence of sarcopenia, being validated for the Brazilian population.¹³

Muscle strength

Muscle strength was assessed with a hand dynamometer for obtaining handgrip strength (HGS). The dynamometers were calibrated before starting data collection. The cut-off points established by Lauretani et al.¹⁴ were used: HGS < 30 kg/f for men and HGS < 20 kg/f for women were considered unfavorable.

Assessment of dietary intake

Dietary intake was estimated by two 24-hour dietary recalls administered on nonconsecutive days. For data to be more valid, multiple-pass recalls were used. To reduce possible reporting errors, utensils (plates, glasses, cups, silverware) were used for amounts. The amounts of foods and preparations were converted to grams or milliliters with a table of cooking measurements.¹⁵ Based on dietary intake data, obtained with the aid of Dietpro®, version 5.5 (Viçosa, Minas Gerais, Brazil), calorie and protein intakes (total and main meals, namely breakfast, lunch, and dinner) were evaluated.

During the fieldwork, dietary recalls were reviewed by nutritionists to identify and resolve any reporting failures. Dietary recalls with a total energy intake lower than 800 kcal or greater than 3,500 kcal were reviewed.

Other variables of interest

A standardized questionnaire was used to characterize the sample in terms of sex, age (in completed years), self-reported ethnicity (Caucasian, Afro-descendant, and Asian), level of education, self-reported comorbidities, and physical activity (weekly frequency).

Statistical analysis

Data were entered into an Excel 2010 spreadsheet with double data entry and analyzed on Stata, version 14 (Stata Corp., College Station, United States). Descriptive statistics, such as mean, standard deviation (SD), and percentage, were used to report the results. For dietary intake variables, which were normally distributed (Kolmogorov-Smirnov test), parametric statistics were used with the performance of analysis of variance (ANOVA test), followed by Tukey test.

Dependent variables (protein from breakfast, lunch, dinner, and protein/kg/day) were associated with independent variables of nutritional status using multiple linear regression

analysis adjusted for sociodemographic variables (gender, age, and education). In all tests, a significance level of 5% or the corresponding p-value was adopted.

Ethical considerations

This study meets the principles of the Brazilian National Health Council Resolution No. 466, of December 12, 2012, and was approved by a Human Research Ethics Committee with opinion No. 1.997.354. All procedures were explained to the patients, and those who agreed to participate signed an informed consent form.

RESULTS

Forty-seven older adults aged between 60 and 96 years were evaluated. Most of them were female (87.23%), widowed (55.32%), and Caucasian (80.85%). Physical activity was reported by 91.49% (Table 1).

Anthropometric data are described in Table 2. The mean BMI was 29.11 kg/m², ranging from 14.15 to 48.06 kg/m². The mean CC was 34.05 (SD = 0.91) cm. The mean WC was 94.52 (SD = 22.73) cm for men and 98.5 (SD = 17.26) cm for women, with no significant difference between sexes ($p = 0.61$). The mean MNA® score was 20.97 (SD = 3.48).

All male participants had reduced muscle strength. Among female participants, 80.0% ($n = 32$) had reduced strength in the left hand and 77.5% ($n = 31$), in the right hand.

The participants reported a mean intake of 1,305.28 kcal/day (19.98% from protein), 65.19 g/day of protein, with 0.93 g protein/kg/day (Table 3). Most older adults reported having three to four meals per day (63.04%), followed by those who had five meals (19.56%), two meals (10.87%), and six meals (6.53%).

In the analysis of the distribution of total and per-meal protein versus the independent variables, protein intake at breakfast was associated with CC ($p = 0.04$) (Table 4). The total amount of protein consumed and its distribution across the other meals (lunch and dinner) were not associated with the analyzed variables.

DISCUSSION

Older adults, mostly Caucasian, female, physically active, at risk of malnutrition, with reduced strength and low level of education, had a mean protein intake lower than the recommended level for preventing muscle loss in this age group. No meal met the recommended protein intake. Lunch had the greatest amount of protein. There was an association between the amount of protein consumed at breakfast and CC.

Protein intake recommendations are traditionally based on studies that estimate the minimal requirement to avoid a negative nitrogen balance. However, these recommendations are currently considered inadequate,⁶ especially for older adults, as they have loss of muscle mass, lower energy intake, lower level of physical activity, and other comorbidities that can affect nitrogen balance and protein anabolism.⁶ The currently

TABLE 1. Sociodemographic, health, and physical activity characteristics of older adults attending a community center.

Variable	% (n)
Sex	
Female	87.23 (41)
Male	12.77 (6)
Age range (years)	
60-69	44.68 (21)
70-79	51.06 (24)
80 or over	4.26 (2)
Race	
Caucasian	80.85 (38)
Afro-descendant	14.89 (7)
Asian	4.26 (2)
Marital status	
Single	2.13 (1)
Married	27.66 (13)
Divorced/separated	14.89 (7)
Widowed	55.32 (26)
Level of education	
Illiterate	21.27 (10)
Primary education*	65.96 (31)
Secondary education*	12.77 (6)
Physical activity	
Yes	91.49 (43)
No	8.51 (4)
Frequency of physical activity (times/week)	
1-2	12.77 (6)
3-4	42.55 (20)
5-7	44.68 (21)
Prevalence of self-reported comorbidities	
Hypertension	39.13 (18)
Diabetes mellitus	26.09 (12)
Osteoporosis	23.91 (11)
Dyslipidemias	28.26 (13)

*Complete or incomplete.

TABLE 2. Descriptive analysis of anthropometric variables of older adults attending a community center.

Variable	Mean	SD	95%CI
Weight (kg)	69.96	15.61	66.76 – 73.17
Height (m)	1.55	0.08	1.54 – 1.57
Body mass index			
LW (%)	4.25	-	-
NW (%)	51.06	-	-
OW (%)	44.69	-	-
RHGS (kg)	16.72	0.60	15.51 – 17.92
LHGS (kg)	17.16	0.66	15.84 – 18.48
SMM (kg)	29.80	0.78	28.25 – 31.36
AC (cm)	31.53	0.79	29.95 – 33.10
Waist circumference (%)			
Low risk	19.15		
Increased risk	17.02		
Greatly increased risk	63.83		
Calf circumference (%)			
Risk of malnutrition	12.77		
Adequate	87.23		
MNA®			
Malnourished (%)	14.89	-	-
Risk of malnutrition (%)	68.08%	-	-
Adequate	17.02%	-	-

SD: standard deviation; CI: confidence interval; LW: low weight; NW: normal weight; OW: overweight. RHGS: right handgrip strength; LHGS: left handgrip strength; SMM: skeletal muscle mass; AC: arm circumference; CC: calf circumference; MNA: Mini-Nutritional Assessment.

TABLE 3. Mean daily and per-meal calorie and protein intake by older adults attending a community center.

Variable	Mean	SD	95%CI
Calorie intake (kcal/day)	1,305.28	495.66	1,202.88 – 1,407.68
Protein intake (g/day)	65.19	39.09	54.94 – 71.43
Protein intake (g/kg/day)	0.93	0.06	0.8137 – 1.0549
Per-meal calorie intake (kcal/%TEI*)			
Breakfast	235.24/18.8	166.20	200.90 – 269.57
Lunch	473.07/36.24	283.19	414.56 – 531.58
Dinner	299.59/22.95	208.49	256.52 – 342.66
Per-meal protein intake (g)			
Breakfast	8.56	0.82	6.92 – 10.19
Lunch	33.85	2.47	28.94 – 38.76
Dinner	19.00	1.73	15.56 – 22.43

*TEI: total energy intake.

TABLE 4. Association of total and per-meal protein intake with the analyzed variables of older adults attending a community center.

Variable	β coefficient	95%CI
Protein/kg/d		
Sex	-0.10	-0.52 – 0.32
RHGS	0.01	-0.02 – 0.03
LHGS	-0.01	-0.04 – 0.01
SMM (kg)	-0.01	-0.03 – 0.01
CP (cm)	-0.01	-0.02 – 0.01
No. of meals	0.06	-0.06 – 0.17
Level of education*	0.18	-0.15 – 0.52
Age†	-0.03	-0.30 – 0.24
Protein at breakfast (g)		
Sex	-3.35	-9.22 – 2.50
RHGS	0.07	-0.33 – 0.49
LHGS	-0.07	-0.46 – 0.31
SMM (kg)	-0.25	-0.56 – 0.04
CC (cm)	0.22	0.01 – 0.43
No. of meals	1.32	-0.29 – 2.93
Level of education*	1.72	-2.89 – 6.73
Age†	-0.67	-3.11 – 4.45
Protein at lunch (g)		
Sex	-5.84	-23.70 – 12.0
RHGS	0.40	-0.85 – 1.67
LHGS	-0.49	-1.67 – 0.69
SMM (kg)	0.12	-0.81 – 1.05
CP (cm)	0.07	-0.58 – 0.72
No. of meals	4.33	-0.59 – 9.27
Level of education*	-0.81	-14.90 – 13.28
Age†	-3.25	-14.8 – 8.30
Protein at dinner (g)		
Sex	9.40	-3.18 – 21.98
RHGS	-0.35	-1.24 – 0.53
LHGS	0.01	-0.83 – 0.83
SMM (kg)	0.36	-0.28 – 1.02
CC (cm)	0.10	-0.36 – 0.55
No. of meals	-0.45	-3.92 – 3.01
Level of education*	6.63	-3.27 – 16.54
Age†	1.43	-6.68 – 9.55

CI: confidence interval; RHGS: right handgrip strength; LHGS: left handgrip strength; SMM: skeletal muscle mass; CC: calf circumference. *Education categorized as illiterate and some level of education. †Age stratified as up to 70 years and 70 years or over.

recommended protein intake is 0.8 g/kg/day.⁸ However, for preventing loss of muscle mass, autonomy, and health with aging, consuming 1.0 to 1.5 g/kg/day of high-quality protein is recommended.⁶

In a randomized, double-blind, placebo-controlled trial of 120 older adults with mild frailty, the participants were supplemented with different dosages (0.8, 1.2, and 1.5 g/kg/d) of whey protein for 12 weeks. A dosage of 1.5 g/kg/day was found to be more efficient to prevent sarcopenia and frailty when compared with the minimum dosage.¹⁶ Another study¹⁷ based on nitrogen balance data showed that consuming 1.2 g/kg/day of protein promotes an adequate balance and is associated with a 40.0% lower loss of muscle mass in older adults when compared with consuming the recommended level (0.8 g/kg/day). Thus, the older adults in our study had a protein intake (0.93 g/kg/d) lower than the recommended level and lower than the level described in other studies.^{5,18} Although no association was observed between total protein intake and the variables of interest, the low protein intake in this sample is consistent with the low muscle strength and the high percentage of those at risk of malnutrition.

The importance of older adults consuming greater amounts of protein is known. However, considerable interest has been recently shown for the pattern of protein distribution across meals, as this seems to be a determining factor for the maintenance of SMM.¹⁹ The explanation is that essential amino acids (EAAs), especially leucine, stimulate MPS,²⁰ and therefore a high-quality protein threshold should be determined for each meal. Also, the EAAs are not stored and, when consumed in excess, are oxidized. Reduced MPS was observed in older adults as a response to a small dose of EAA, but the response to a higher intake of EAA, corresponding to 25-30 g of protein, was normal.²¹

Young people need 24 g/meal of protein, while older adults need 40 g/meal to achieve maximal MPS.⁷ Thus, aging is associated with a change in the dose-response curve for protein intake and MPS, and higher doses are required to maximally stimulate MPS.⁷ Based on these values, the protein intake found in our study (65.19 g/d) would be sufficient to meet the ideal recommendation in only one of the three main meals. However, the recommendation of 40 g/meal was met in none of the meals. Lunch was the meal with the highest protein intake (33.85 [SD = 2.47 g]). A similar study conducted in Mexico²² also described a higher protein intake at lunch. However, the pattern of protein distribution across meals remains controversial. Findings from other countries described dinner as the meal with the highest protein intake, accounting for 40-50% of total protein intake.^{5,23} However, despite specific cultural differences between countries, it

seems that the time of day with the highest protein intake does not influence the association between adequate dietary protein intake and disabilities.²²

Breakfast is traditionally the meal with the lowest protein intake,^{4,19} as observed in our study (8.56 g). Higher values were described for older Mexicans. Those who consumed 30 g of protein at one of the daily meals (as in our study) consumed 18.7 g of protein at breakfast. Older Norwegians, in turn, consumed 17.0 g (SD = 8.2) of protein at breakfast, which is usually very similar to lunch (they eat bread with some kind of spread).⁵ The same was observed in the Netherlands, where 10.3 to 12.3 g of protein were described at breakfast,²³ with bread, milk, and milk products being the main sources of dietary protein.²⁴ Dietary intake data on older Brazilians²⁵ demonstrated that butter/margarine, bread, coffee, sugar, and milk are the foods traditionally present in the breakfast meal of this population, or they are meal substitutes, especially at dinner.

Low protein intake at breakfast was significantly associated with CC in this sample. CC is a measure that reflects protein reserve and is considered a marker of SMM.²⁶ Reduced SMM is directly associated with a higher risk of sarcopenia.³ Thus, our findings suggest that the lower amount of protein at breakfast is associated with a higher risk of sarcopenia and frailty. The total amount of protein consumed and the protein content in other meals had no association with the analyzed variables. Bollwein et al.¹⁸ described a lower protein intake at breakfast and a higher intake at lunch among frail older Germans. In older Mexicans without limitations, consuming two or three meals/day with 30 g of protein each was associated with a lower risk of physical disability.²²

These findings are important for public health and for the management of care for older adults, as low total protein intake and inadequate protein distribution across meals (40 g/meal) may lead to loss of muscle mass and increased risk of sarcopenia.^{4,6,7} In addition to quantity, protein quality influences MPS.⁶ Protein quality is characterized by its amino acid composition and digestion and absorption factors. Animal protein (milk, meat, and eggs) seems to contain a higher amount of EAA and thus promotes increased MPS when compared with plant protein (soybean, wheat).²⁷ Therefore, given the low dietary protein intake, especially at breakfast, observed in older Brazilians, encouraging the inclusion of animal-based protein sources, such as eggs, milk, and milk products, is an important strategy to improve SMM synthesis and to prevent the development of sarcopenia and fragility.

Furthermore, energy intake is a key factor when evaluating protein intake. The mean energy intake of the sample was lower than that obtained for the older Brazilian population according to 2017–2018 Consumer Expenditure Survey (Pesquisa de Orçamento Familiar, POF) data, which described the consumption of 1,709 kcal/day among older men and 1,409 kcal/day among older women.²⁸ A lower energy intake can be determining for a lower protein intake; however, despite the low calories in this sample, the percentage of protein energy (19.98%) is consistent with that recommended in the literature⁸ and that described for older Brazilians (19.0% among men and 17.9% among women).²⁸ In addition to energy, factors such as anorexia, changes in food preferences, food insecurity, food culture, difficulty preparing food, and dental and swallowing problems may lead to a low protein intake in this population.^{29,30}

It is worth noting that this study used standardized protocols and validated tools to obtain all the reported data. As national data on protein intake, protein distribution across meals, and the relationship of protein distribution with the health of older adults are limited, the relevance of this study lies on widening the discussion of the topic. The results should encourage the development of new population-based studies to better understand the effects of protein distribution across meals on the muscle mass of older adults.

Although this study makes important contributions to the medical literature, it has some limitations. The first is the sample size, considering that the number of older adults who fully completed the study protocol was low. The second is the predominance of women in the sample, given the low level of participation of older men. Another limitation is that only older adults who attended the community center participated in the investigation; thus, the sample is not representative of older adults living in the community. Therefore, caution should be exercised when applying these results to other populations.

CONCLUSION

The older adults in the study sample had low total and per-meal protein intakes. Breakfast was the meal that provided the lowest protein intake, and this was associated with CC, a marker for the diagnosis of sarcopenia in older adults. Encouraging the consumption of protein-rich foods at breakfast, preferably animal protein, is a relevant dietary strategy for preventing and treating sarcopenia and frailty in this population.

Conflict of interest

The authors declare no conflicts of interest.

Funding

This study received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Authors' contribution

LBS: project administration, conceptualization, data curation, writing – original draft, writing – review & editing, investigation, methodology, supervision. KAM: conceptualization, data curation, writing – review & editing. RAB: formal analysis, data curation, writing – original draft, writing – review & editing.

REFERENCES

1. Instituto Brasileiro de Geografia e Estatística. Projeção da população do Brasil e das Unidades da Federação. Available from: <https://www.ibge.gov.br/apps/populacao/projecao/index.html>. Accessed in Mar 28, 2021.
2. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in older people. *Age Ageing*. 2010;39(4):412-23. <http://doi.org/10.1093/ageing/afq034>.
3. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*. 2019;48(1):16-31. <http://doi.org/10.1093/ageing/afy169>.
4. Traylor DA, Gorissen SHM, Phillips SM. Perspective: protein requirements and optimal intakes in aging: are we ready to recommend more than the recommended daily allowance? *Adv Nutr*. 2018;9(3):171-82. <http://doi.org/10.1093/advances/nmy003>.
5. Nygård LK, Dahl L, Mundal I, Benth SJ, Rokstad AMM. Protein intake, protein mealtime distribution and seafood consumption in elderly norwegians: associations with physical function and strength. *Geriatrics (Basel)*. 2020;5(4):100. <http://doi.org/10.3390/geriatrics5040100>.
6. Phillips SM, Chevalier S, Leidy HJ. Protein "requirements" beyond the RDA: implications for optimizing health. *Appl Physiol Nutr Metab*. 2016;41(5):565-72. <http://doi.org/10.1139/apnm-2015-0550>.
7. Moore DR, Churchward-Venne TA, Witard O, Breen L, Burd NA, Tipton KD, et al. Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men. *J Gerontol A Biol Sci Med Sci*. 2015;70(1):57-62. <http://doi.org/10.1093/gerona/glu103>.
8. Trumbo P, Schlicker S, Yates AA, Poos M, Food and Nutrition Board of the Institute of Medicine, The National Academies. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein and amino acids. *J Am Diet Assoc*. 2002;102(11):1621-30. [http://doi.org/10.1016/s00028223\(02\)90346-9](http://doi.org/10.1016/s00028223(02)90346-9).
9. Lohman TG, Roche AF, Martorell R. Anthropometric standardization reference manual. Champaign: Human Kinetics Books; 1988.
10. Lipschitz DA. Screening for nutritional status in the elderly. *Prim Care*. 1994;21(1):55-67. PMID: 8197257.
11. National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation*. 2002;106(25):3143-421. PMID: 12485966.
12. Lee RC, Wang Z, Heo M, Ross R, Janssen I, Heymsfield SB. Total-body skeletal muscle mass: development and cross-validation of anthropometric prediction models. *Am J Clin Nutr*. 2000;72(3):796-803. <http://doi.org/10.1093/ajcn/72.3.796>.
13. Rech CR, Dellagrana RA, Marucci MFN, Petroski EL. Validity of anthropometric equations for the estimative of muscle mass in elderly. *Rev Bras Cineantropom Desempenho Hum*. 2012;14(1):23-31. <http://doi.org/10.5007/1980-0037.2012v14n1p23>.
14. Lauretani F, Russo CR, Bandinelli S, Bartali B, Cavazzini C, Di Iorio A, et al. Age-associated changes in skeletal muscles and their effect on mobility: an operational diagnosis of sarcopenia. *J Appl Physiol* (1985). 2003;95(5):1851-60. <http://doi.org/10.1152/japplphysiol.00246.2003>.
15. Pinheiro ABV, Lacerda EMA, Benzecty EH, Gomes MCS, Costa VM. Tabela para Avaliação de Consumo Alimentar em Medidas Caseiras. 5th. ed. Rio de Janeiro: Atheneu; 2008.
16. Park SW, Goodpaster BH, Lee JS, Kuller LH, Boudreau R, Rekenneir N, et al. Excessive loss of skeletal muscle mass in older adults with type 2 diabetes. *Diabetes Care*. 2009;32(11):1993-7. <http://doi.org/10.2337/dc09-0264>.
17. Houston DK, Nicklas BJ, Ding J, Harris TB, Tylavsky FA, Newman AB, et al. Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the Health, Aging, and Body Composition (Health ABC) Study. *Am J Clin Nutr*. 2008;87(1):150-5. <http://doi.org/10.1093/ajcn/87.1.150>.
18. Bollwein J, Diekmann R, Kaiser MJ, Bauer JM, Uter W, Sieber CC, et al. Distribution but not amount of protein intake is associated with frailty: a cross-sectional investigation in the region of Nürnberg. *Nutr J*. 2013;12:109. <http://doi.org/10.1186/1475-2891-12-109>.
19. Farsijani S, Morais JA, Payette H, Gaudreau P, Shatenstein B, Gray-Donald K, et al. Relation between mealtime distribution of protein intake and lean mass loss in free-living older adults of the NuAge study. *Am J Clin Nutr*. 2016;104(3):694-703. <http://doi.org/10.3945/ajcn.116.130716>.
20. Santos CS, Nascimento FEL. Consumo isolado de aminoácidos de cadeia ramificada e síntese de proteína muscular em humanos: uma revisão bioquímica. *Einstein (São Paulo)*. 2019;17(3):eRB4898. https://doi.org/10.31744/einstein_journal/2019RB4898.
21. Paddon-Jones D, Sheffield-Moore M, Zhang XJ, Volpi E, Wolf SE, Aarsland A, et al. Amino acid ingestion improves muscle protein synthesis in the young and elderly. *Am J Physiol Endocrinol Metab*. 2004;286(3):E321-8. <http://doi.org/10.1152/ajpendo.00368.2003>.
22. Gaytán-González A, Ocampo-Alfaro MJ, Torres-Naranjo F, Arroniz-Rivera M, González-Mendoza RG, Gil-Barreiro M, et al. The consumption of two or three meals per day with adequate protein content is associated with lower risk of physical disability in Mexican adults aged 60 years and older. *Geriatrics (Basel)*. 2020;5(1):1. <http://doi.org/10.3390/geriatrics5010001>.
23. Tieland M, Borgonjen-Van den Berg KJ, Van Loon LJC, Groot LCPGM. Dietary protein intake in Dutch elderly people: a focus on protein sources. *Nutrients*. 2015;7(12):9697-706. <http://doi.org/10.3390/nu7125496>.
24. Tieland M, Borgonjen-Van den Berg KJ, van Loon LJC, Groot LCPGM. Dietary protein intake in community-dwelling, frail, and institutionalized elderly people: scope for improvement. *Eur J Nutr*. 2012;51(2):173-9. <http://doi.org/10.1007/s00394-011-0203-6>.
25. Ferreira MPN, Previdelli AN, Freitas TI, Marques KM, Goulart RMM, Aquino RC. Padrões dietéticos e fatores associados em idosos. *Rev Bras Geriatr Gerontol*. 2017;20(4):538-49. <https://doi.org/10.1590/1981-22562017020.160115>.
26. Kawakami R, Murakami H, Sanada K, Tanaka N, Sawada SS, Tabata I, et al. calf circumference as a surrogate marker of muscle mass for diagnosing sarcopenia

- in Japanese men and women. *Geriatr Gerontol Int.* 2015;15(8):969-76. <http://doi.org/10.1111/ggi.12377>.
27. van Vliet S, Burd NA, van Loon LJC. The skeletal muscle anabolic response to plant- versus animal-based protein consumption. *J Nutr.* 2015;145(9):1981-91. <http://doi.org/10.3945/jn.114.204305>.
28. Instituto Brasileiro de Geografia e Estatística. Pesquisa de orçamentos familiares 2017-2018: análise do consumo alimentar pessoal no Brasil. Rio de Janeiro: IBGE; 2020.
29. Bomfim RA, Souza LB, Corrente JE. Tooth loss and its relationship with protein intake by elderly Brazilians-a structural equation modelling approach. *Gerodontology.* 2018;35(1):51-8. <http://doi.org/10.1111/ger.12317>.
30. Valenzuela RER, Ponce JA, Morales-Figueroa GG, Muro KA, Carreón VR, Alemán-Mateo H. Insufficient amounts and inadequate distribution of dietary protein intake in apparently healthy older adults in a developing country: implications for dietary strategies to prevent sarcopenia. *Clin Interv Aging.* 2013;8:1143-8. <http://doi.org/10.2147/CIA.S49810>.